


Synonym

PDCD1, PD1, CD279, SLEB2

Source

Biotinylated Human PD-1, Avitag, His Tag (recommended for biopanning) (PD1-H82E4) is expressed from human 293 cells (HEK293). It contains AA Leu 25 - Gln 167 (Accession # Q15116-1).

N-terminal Sequence Analysis: Leu 25

Molecular Characterization


PD-1 (Leu 25 - Gln 167)
Q15116-1

This protein carries an Avi tag (Avitag™) at the C-terminus, followed by a polyhistidine tag.

The protein has a calculated MW of 18.6 kDa. The protein migrates as 35-45 kDa under reducing (R) condition (SDS-PAGE) due to glycosylation.

Application

PD1-H82E4 works best for experiments that test the binding between PD-1 and candidate antibodies, such as biopanning and other relevant assays.

This product is NOT suitable for testing PD1-PDL1 binding. For this type of application, we strongly recommend you to choose PD1-H82F2 as an alternative.

Biotinylation

Biotinylation of this product is performed using Avitag™ technology. Briefly, the single lysine residue in the Avitag is enzymatically labeled with biotin.

Biotin:Protein Ratio

The biotin to protein ratio is *0.5-1* as determined by the HABA assay.

Endotoxin

Less than 1.0 EU per µg by the LAL method.

Purity

>95% as determined by SDS-PAGE.

Formulation

Lyophilized from 0.22 µm filtered solution in PBS, pH7.4. Normally trehalose is added as protectant before lyophilization.

Contact us for customized product form or formulation.

Reconstitution

Please see Certificate of Analysis for specific instructions.

For best performance, we strongly recommend you to follow the reconstitution protocol provided in the CoA.


Storage

For long term storage, the product should be stored at lyophilized state at -20°C or lower.

Please avoid repeated freeze-thaw cycles.


No activity loss was observed after storage at:

- 4-8°C for 12 months in lyophilized state;
- -70°C for 3 months under sterile conditions after reconstitution.

SDS-PAGE

Biotinylated Human PD-1, Avitag, His Tag (recommended for biopanning) on SDS-PAGE under reducing (R) condition. The gel was stained overnight with Coomassie Blue. The purity of the protein is greater than 95%.

Bioactivity-ELISA



Immobilized Nivolumab at 2 µg/mL (100 µL/well) can bind Biotinylated Human PD-1, Avitag, His Tag (recommended for biopanning) (Cat. No. [PD1-H82E4](#)) with a linear range of 0.8-13 ng/mL (QC tested).

Background

Programmed cell death protein 1 (PD-1) is also known as CD279 and PDCD1, is a type I membrane protein and is a member of the extended CD28/CTLA-4 family of T cell regulators. PDCD1 is expressed on the surface of activated T cells, B cells, macrophages, myeloid cells and a subset of thymocytes. PD-1 has two ligands, PD-L1 and PD-L2, which are members of the B7 family. PD-L1 is expressed on almost all murine tumor cell lines, including PA1 myeloma, P815 mastocytoma, and B16 melanoma upon treatment with IFN-γ. PD-L2 expression is more restricted and is expressed mainly by DCs and a few tumor lines. PD1 inhibits the T-cell proliferation and production of related cytokines including IL-1, IL-4, IL-10 and IFN-γ by suppressing the activation and transduction of PI3K/AKT pathway. In addition, coligation of PD1 inhibits BCR-mediated signal by dephosphorylating key signal transducer. In vitro, treatment of anti-CD3 stimulated T cells with PD-L1-Ig results in reduced T cell proliferation and IFN-γ secretion. Monoclonal antibodies targeting PD-1 that boost the immune system are being developed for the treatment of cancer.

References

- (1) [Ishida Y., et al., 1992, EMBO J. 11 \(11\): 3887-95.](#)
- (2) [Blank C., et al., 2007, Cancer Immunol. Immunother. 56 \(5\): 739-45.](#)
- (3) [Agata Y., et al., 1996, Int. Immunol. 8 \(5\): 765-72.](#)
- (4) [Freeman GJ., et al., 2000, J. Exp. Med. 192 \(7\): 1027-34.](#)
- (5) [Latchman Y., et al., 2001, Nat. Immunol. 2 \(3\): 261-8.](#)
- (6) [Yamazaki T., et al., 2002, J. Immunol. 169 \(10\): 5538-45.](#)

Please contact us via TechSupport@acrobiosystems.com if you have any question on this product.